1 **Explicative note to this paper**

2 This paper was submitted to **BAMS** on June 27, 2012, it was accepted for
3 publication by editor C. Deser on Dec. 27, 2012 and retracted by the senior **BAMS** editor
4 J. Rosenfeld on Oct. 24, 2013. The reason given was that a much shorter review of
5 similar material (29% of the length) had been published in the weekly magazine **EOS**
6 ([Lovejoy, S., 2013: What is Climate?: EOS, 94, No. 1, 1 January 2013, p1-2.](#)). Lovejoy
7 was therefore accused of attempting “concurrent dual publication”.

8 The background was that at the moment of submission, the draft paper was posted
9 on Lovejoy’s personal web site. It rapidly gained notoriety in the blogosphere and came
10 to the attention of the **EOS** editors R. Pielke Sr. and B. Richman. In August 2012, this
11 much shorter review was explicitly commissioned by **EOS** to cover similar material but
12 in much shorter form and for a quite a different audience. The practice of **EOS**
13 commissioning short reviews of new work covered at greater length in other journals is
14 common and there was certainly no attempt at dual publication. More details on this
15 unfortunate episode may be found at in the chronology document
16 ([http://www.physics.mcgill.ca/~gang/eprints/eprintLovejoy/neweprint/ChronologyBAMS](http://www.physics.mcgill.ca/~gang/eprints/eprintLovejoy/neweprint/ChronologyBAMS.EOS.25.10.13.pdf)
17 [.EOS.25.10.13.pdf](http://www.physics.mcgill.ca/~gang/eprints/eprintLovejoy/neweprint/ChronologyBAMS.EOS.25.10.13.pdf)).
18

The climate is not what you expect

S. Lovejoy¹, D. Schertzer²

¹Physics, McGill University

3600 University st.,

Montreal, Que., Canada

Email: lovejoy@physics.mcgill.ca

²CEREVE

Université Paris Est,

6-8, avenue Blaise Pascal

Cité Descartes

77455 MARNE-LA-VALLEE Cedex, France

The climate is not what you expect

Capsule Summary:

Contrary to popular ideas about the climate, what you really expect is macroweather. On scales of a human generation, the climate is what you get.

Abstract:

Prevailing definitions of climate are not much different from “the climate is what you expect, the weather is what you get”. Using a variety of sources including reanalyses and paleo data, and aided by notions and analysis techniques from Nonlinear Geophysics, we argue that this dictum is fundamentally wrong. In addition to the weather and climate, there is a qualitatively distinct intermediate regime extending over a factor of ≈ 1000 in scale. For example, mean temperature fluctuations increase up to about 5 K at 10 days (the lifetime of planetary structures), then decrease to about 0.2 K at 30 years, and then increase again to about 5 K at glacial-interglacial scales.

Both deterministic GCM’s with fixed forcings (“control runs”) and stochastic turbulence-based models reproduce the first two regimes, but not the third. The middle regime is thus a kind of “macroweather” not “high frequency climate”. Averaging macroweather over periods increasing to ≈ 30 yrs yields apparently converging values: macroweather is “what you expect”. Macroweather averages over ≈ 30 years have the lowest variability, they yield well defined “climate states” and justify the otherwise ad hoc “climate normal” period. However, moving to

longer periods, these states increasingly fluctuate: just as with the weather, the climate changes in an apparently unstable manner; the climate is *not* what you expect. Similarly, we may categorize climate forcings according to whether their fluctuations decrease or increase with scale and this has important implications for GCM's and for climate change and climate predictions.

1. Introduction

In his monumental “Climate: Past, Present, and Future” Horace Lamb argued that the early scientific view was “climate as constant” [Lamb, 1972]. Reflecting this, in 1935 the International Meteorological Organization adopted 1901-1930 as the “climatic normal period”. Following the post war cooling, this view evolved: for example the official American Meteorological Society glossary [Huschke, 1959] defined the climate as “the synthesis of the weather” and then “...the climate of a specified area is represented by the statistical collective of its weather conditions during a specified interval of time (usually several decades)”. Although this new definition in principle allows for climate change, the period 1931-1960 soon became the new “normal”, the ad hoc 30 year duration became entrenched, today 1961-1990 is commonly used. Mindful of the extremes, Lamb warned against reducing the climate to just “average weather”, while viewing the climate as “...the sum total of the weather experienced at a place in the course of the year and over the years”, [Lamb, 1972].

Lamb's essentially modern view allows for the possibility of climate change and is closely captured by the popular expression: “The climate is what you expect, the weather is what you get” (the character Lazurus Long in [Heinlein, 1973], but often attributed to Mark Twain). It is also close to the US National Academy of

77 Science definition: “Climate is conventionally defined as the long-term statistics of
78 the weather...” [*Committee on Radiative Forcing Effects on Climate*, 2005] which
79 improves on the “the climate is what you expect” idea only a little by proposing:
80 “...to expand the definition of climate to encompass the oceanic and terrestrial
81 spheres as well as chemical components of the atmosphere”.

82 The Twain/Heinlein expression was strongly endorsed by the late E. Lorenz
83 who stated: “Before embarking on a search for an ideal definition (of climate)
84 assuming one exists, let me express my conviction that such a definition, when
85 found must agree in spirit with the statement, “climate is what you expect”.” [*Lorenz*,
86 1995]. He then proposed several definitions based on dynamical systems theory
87 and strange attractors (see also [*Palmer*, 2005]).

88 A variant on this, motivated by GCM modeling, was proposed by [*Bryson*, 1997]
89 (criticized by [*Pielke*, 1998]): “Climate is the thermodynamic/ hydrodynamic status of
90 the global boundary conditions that determine the concurrent array of weather patterns.”
91 He explains that whereas “weather forecasting is usually treated as an *initial value*
92 problem ... climatology deals primarily with a *boundary condition* problem and the
93 patterns and climate devolving there from”. This definition could be paraphrased “for
94 given boundary conditions, the climate is what you expect”. This and similar views
95 provide the underpinnings for much of current climate prediction, including the recent
96 idea of “seamless forecasting” (e.g. [*Palmer et al.*, 2008], [*Palmer*, 2012]) in which
97 seasonal scale model validation is applied to climate scale predictions (for a recent
98 discussion, see [*Pielke et al.*, 2012]).

99 There are two basic problems with the Twain/Heinlein dictum and its variants.

100 The first is that they are based on an abstract weather - climate dichotomy, they are not
101 informed by empirical evidence. The glaring question of how long is “long” is either
102 decided subjectively or taken by fiat as the WMO’s “normal” 30 year period. The second
103 problem – that will be evident momentarily – is that it assumes that the climate is nothing
104 more than the long-term statistics of weather. If we accept the usual interpretation – that
105 we “expect” averages - then the dictum means that averaging weather over long enough
106 time periods converges to the climate. With regards to external forcings, one could argue
107 that this notion could still implicitly include the atmospheric response i.e. with averages
108 converging to slowly varying “responses”. However, it implausibly excludes the
109 appearance of any new “slow”, internal climate processes leading to fluctuations growing
110 rather than diminishing with scale.

111 To overcome this objection, one might adopt a more abstract interpretation of
112 what we “expect”. For example if the climate is defined as the probability distribution of
113 weather states, then all we “expect” is a random sample. However even this works only
114 inasmuch as it is possible to merge fast weather processes with slow climate processes
115 into a single process with a well defined probability distribution. While this may satisfy
116 the theoretician, it is unlikely to impress the layman. This is particularly true since to be
117 realistic we will see that one of the regimes in this composite model must have the
118 property that fluctuations converge whereas in the other, they diverge with scale. To use
119 the single term “climate” to encompass the two opposed regimes therefore seems at best
120 unhelpful and at worst misleading.

121 The main purpose of this paper is show that the weather-climate dichotomy is
122 empirically untenable, that hiding between the two is a missing middle range spanning a

factor of a thousand in scale (≈ 10 days to ≈ 30 -100 years) characterized by qualitatively different dynamics. This new low frequency weather regime was dubbed “macroweather” since it is a kind of large scale weather (not small scale climate) regime [Lovejoy and Schertzer, 2013], it fundamentally clarifies the distinction between weather and climate, the status and role of GCM models and the notion of climate predictability.

2. The variability characterized by spectral composites

Notwithstanding the existence of several strong periodicities (notably diurnal and annual), the atmosphere is highly variable over huge ranges of space-time scales. In addition, it has long been recognized (e.g. [Lovejoy and Schertzer, 1986], [Wunsch, 2003]) that even at the longer climate scales, the contribution to the variability from specific frequencies associated with specific quasi periodic processes is small: the great majority of the variance in the spectrum is from the continuous background “noise”. Any objectively based definition of weather or climate must therefore start from a clear picture of the atmosphere’s temporal variability over wide scale ranges.

The first - and still most ambitious - single composite spectrum of atmospheric variability [Mitchell, 1976], ranged from hours to the age of the earth ($\approx 10^{-4}$ to 10^{10} yrs), fig. 1 shows a modern version. Given the rudimentary quality of the data at that time, Mitchell admitted that his composite was mostly an “educated guess” (indeed, over the range $\approx 10^4$ to $\approx 10^8$ yrs, [Shackleton and Imbrie, 1990] empirically found that Mitchell had underestimated the variation of the spectral density by a factor of $\approx 10^5$!). His framework reflected the prevailing idea that there

were numerous roughly periodic processes, superposed onto a continuous background spectrum $E(\omega)$ made up of a hierarchy of white noise processes and their integrals (i.e. Ornstein-Uhlenbeck processes with spectra $E(\omega) \approx \omega^{-\beta}$ with $\beta = 0, 2$ respectively where β is the spectral exponent: the negative slope on the log-log plot in fig. 1). The spectral spikes were therefore superposed on a spectrum consisting of a series of “shelves” and represented distinct physical processes. Mitchell explained his idea as follows:

“As we scan the spectrum from the short-wave end toward the longer wave regions, at each point where we pass through a region of the spectrum corresponding to the time constant of a process that adds variance to the climate, the amplitude of the spectrum increases by a constant increment across all substantially longer wavelengths. In other words, each stochastic process adds a shelf to the spectrum at an appropriate wavelength” [Mitchell, 1976].

By the early 1980’s, following the explosion of scaling (fractal) ideas it was realized that scale invariance was a very general symmetry principle often respected by nonlinear dynamics, including many geophysical processes and turbulence. The signature of a scaling process is a power law spectrum, linear on a log-log plot. Although in order to accommodate the wide range of scales, Mitchell had found it “necessary to resort to logarithmic coordinates”, there was no implication that the underlying processes might have nontrivial scaling over any significant range. In contrast, scaling symmetries, were explicitly invoked to justify

the alternative composite picture ([*Lovejoy and Schertzer, 1984; Lovejoy and Schertzer, 1986*] which profited from early ocean and ice core paleotemperatures. These analyses already clarified the following points: a) the distinction between the variability of regional and global scale temperatures, b) that there was a scaling range for global averages between scales of the order of 10 yrs (τ_c in the notation here) up to $\approx 40 - 50$ kyrs with an exponent $\beta_c \approx 1.8$, c) that a scaling regime with this exponent could quantitatively explain the magnitudes of the temperature swings between interglacials (“ice ages”): the “interglacial window” (see below).

In the last 15 years this picture has been supported by the quite similar scaling composites of [*Pelletier, 1998*] and [*Huybers and Curry, 2006*]. The latter in particular made a data intensive study of the scaling of many different types of paleotemperatures collectively spanning the range of about 1 month to nearly 10^6 years. In addition, even without producing composites, other authors shared the scaling framework, e.g. [*Koscielny-Bunde et al., 1998*], [*Talkner and Weber, 2000*], [*Ashkenazy et al., 2003; Rybski et al., 2008*]. Their results are qualitatively very similar - including the positions of the scale breaks; the main innovations are a) the increased precision on the β estimates and b) the basic distinction made between continental and oceanic spectra including their exponents. We could also mention the composite of [*Fraedrich et al., 2009*] which is a modest adaptation of Mitchell’s: it innovates by introducing a single scaling regime from ≈ 3 to ≈ 100 yrs.

Using real temperature and paleotemperature data, examples showing the behaviors in the three different regimes are graphically illustrated in fig. 2. Notice that in the weather regime (bottom) the temperature seems to “wander” up or down

like a drunkard's walk so that temperature differences typically increase over longer and longer periods. Turning to the middle macroweather series, we see that it has a totally different appearance with successive fluctuations on the contrary tending to cancel each other out, i.e. with decreases followed by partially cancelling increases (and visa versa). At first sight, this vindicates the "climate is what you get" idea since averages over longer and longer periods will clearly converge. From this, we anticipate that at decadal - or certainly at centennial scales - that we will see at most smooth, slow variations. However, when we turn to the century resolution climate series (top) on the contrary, we once again see weather-like wandering.

Below we shall see that the fluctuations over a time interval Δt are in each case roughly scaling (power laws) of the form Δt^H so that the sign of H qualitatively distinguishes the "wandering" ($H>0$) or "cancelling" ($H<0$) behaviors (since $\beta \approx 2H+1$, the critical value $H = 0$ roughly corresponds to a critical spectral exponent $\beta = 1$). In nonlinear dynamical systems, power laws arise when over a range of scales there are no processes strong enough to break the scaling symmetry. Another way of putting this is to say that the dominant dynamical processes occur in synergy over a wide range of scales, with the resulting behavior displaying no characteristic size or duration.

3. Evidence for scaling in the three regimes

Taken individually, for the weather ($\Delta t \approx \tau_w$; $\tau_w \approx 10$ days), macroweather ($\tau_w < \Delta t < \tau_c$; $\tau_c \approx 30 - 100$ yrs), and climate ($\Delta t > \tau_c$), there are now many studies supporting the scaling picture and estimating various scaling exponents in each.

Starting with the climate regime, numerous paleo temperature series (mostly from ice and ocean cores) have been analyzed and there is broad agreement on their scaling nature with spectral exponents estimated in the range $\beta_c \approx 1.3$ to 2.1 over range from hundreds to tens of thousands of years, [Lovejoy and Schertzer, 1986], [Schmitt *et al.*, 1995], [Ditlevsen *et al.*, 1996], [Pelletier, 1998], [Ashkenazy *et al.*, 2003], [Wunsch, 2003], [Huybers and Curry, 2006], [Blender *et al.*, 2006], [Lovejoy and Schertzer, 2012c]. These analyses employed diverse techniques including spectra, difference and Haar structure functions as well as Detrended Fluctuation Analysis so that the results are fairly robust. In addition, as discussed below (fig. 3), further analyses from surface temperatures, multiproxy reconstructions and 138 year long Twentieth Century reanalysis (20CR), lend this further quantitative support.

Similarly, in the macroweather regime, there are now many studies finding scaling with spectral exponents $\beta_{mw} < 1$, e.g. for the temperature; with some variation in β_{mw} between oceans and continents, northern latitudes and tropics: [Lovejoy and Schertzer, 1986], [Pelletier, 1998], [Huybers and Curry, 2006], [Fraedrich and Blender, 2003], Koscielny-Bunde *et al.*, 1998, Bunde *et al.*, 2004, [Eichner *et al.*, 2003], [Lennartz and Bunde, 2009], [Blender *et al.*, 2006], [Fraedrich *et al.*, 2009], [Lanfredi *et al.*, 2009]. Since β_{mw} is small, log-log spectra appear as fairly flat hence the original term “spectral plateau”. A review of the ubiquitous empirical evidence for this include analyses of the temperature, wind, humidity, geopotential height, rain, vertical wind, and the North Atlantic Oscillation, Southern Oscillation and Pacific Decadal Oscillation indices [Lovejoy and Schertzer, 2010], [Lovejoy and Schertzer, 2013].

236 Of the three regimes, the only one where the idea of a roughly scaling
 237 background spectrum is still somewhat controversial is the higher frequency
 238 weather regime (scales $< \tau_w \approx 10$ days). To understand the debate, recall that the
 239 classical turbulence theories describing the statistical variability in the weather
 240 regime are all based on isotropic scaling, the most famous being the Kolmogorov k
 241 $^{5/3}$ spectrum for the wind (k is a wavenumber). However, the strong vertical
 242 atmospheric stratification prevents isotropic scaling from holding over any scale
 243 ranges spanning the scale thickness of the atmosphere (≈ 10 km). One must
 244 therefore abandon either the scaling or the isotropy assumption. Following
 245 Kraichnan's development of 2-D turbulence and Charney's extension to (still
 246 essentially 2D) "quasi geostrophic" turbulence, the usual choice was to retain
 247 isotropy and to divide the dynamics into 2D isotropic (large scale) and 3D isotropic
 248 (small) scale regimes ([*Kraichnan*, 1967], [*Charney*, 1971]). However, starting with
 249 [*Schertzer and Lovejoy*, 1985], a growing body of evidence and theory has
 250 supported the alternative anisotropic scaling hypothesis. Thanks both to modern
 251 empirical evidence (e.g. the review [*Lovejoy and Schertzer*, 2010], [*Lovejoy and*
 252 *Schertzer*, 2013] and a recent massive aircraft study [*Pinel et al.*, 2012]), but also to
 253 theoretical arguments showing that the governing equations are symmetric with
 254 respect to anisotropic scaling symmetries ([*Schertzer et al.*, 2012]), the question
 255 increasingly has been settled in favor of anisotropic scaling (see the recent debate
 256 [*Lovejoy et al.*, 2009], [*Lindborg et al.*, 2010a; *Lindborg et al.*, 2010b], [*Lovejoy et*
 257 *al.*, 2010], [*Schertzer et al.*, 2011], [*Yano*, 2009], [*Schertzer et al.*, 2012]). The

implications of this anisotropic spatial scaling for the temporal statistics are discussed in [Radkevitch *et al.*, 2008] and [Lovejoy and Schertzer, 2010].

A review of diverse evidence from reanalyses, in situ and remotely sensed data ([Lovejoy and Schertzer, 2010], [Lovejoy and Schertzer, 2013]) shows that for wind, temperature, humidity, pressure, short and long wave radiances, β_w is commonly in the range 1.5 -2 (certainly >1 , hence $H>0$). The existence of a basic transition in the range $\approx 5 - 20$ days has been recognized at least since [Van der Hoven, 1957] who noted a low frequency spectral “bump” at around 5 days. Later, the corresponding features in the temperature and pressure spectra were termed “synoptic maxima” by [Kolesnikov and Monin, 1965] and [Panofsky, 1969]. More recently, in the same spirit as Mitchell, the transition has been modeled (e.g. [AchutaRao and Sperber, 2006]) as an Ornstein-Uhlenbeck process i.e. with $\beta_w = 2$, $\beta_{mw} = 0$, corresponding to $H_w = 1/2$, $H_{mw} = -1/2$, although as can be seen in fig. 3 (discussed below), this is not a very accurate approximation and can be misleading. Finally, [Vallis, 2010] proposed a (nonscaling) mechanism by suggesting that τ_w is determined by the lifetimes of baroclinic instabilities. These were estimated by the inverse Eady growth rate (τ_{Eady}) which yielded $\tau_{Eady} \approx 4$ days. However this result requires a linearization of the equations about a hypothetical state having uniform shear and uniform stratification across the entire troposphere. In comparison, the real troposphere has highly nonuniform shears and stratifications, its variability is so strong that it is characterized by anomalous scaling exponents throughout [Lovejoy *et al.*, 2007] (including $H \approx 0.75$ for the horizontal wind in the vertical direction). Another difficulty with using $\tau_{Eady} \approx \tau_w$ is that τ_{Eady} is inversely proportional to the

Coriolis parameter so that it diverges at the equator whereas the empirical τ_w is not very sensitive to latitude.

Indeed, a seductive feature of the (anisotropic) scaling framework is that it fairly accurately predicts the weather to macroweather transition scale $\tau_w \approx 10$ days. The argument is as follows: the sun provides $\approx 240 \text{ W/m}^2$ of heating with a 2% efficiency of conversion to kinetic energy ([*Monin, 1972*]). The energy is distributed reasonably uniformly over the troposphere, leading to a turbulent energy flux density (ϵ) close to the observed global value $\epsilon \approx 10^{-3} \text{ W/kg}$ ([*Lovejoy and Schertzer, 2010*]; this is the flux of energy from large to small scales). The model predicts that this turbulent energy flux is the fundamental driver of the horizontal dynamics and thus that planetary structures have eddy-turnover times of $\approx \epsilon^{-1/3} L_e^{2/3} \approx 10$ days where $L_e = 20000 \text{ km}$ is the largest great circle distance on the earth. The analogous calculation for the ocean using the empirical ocean turbulent flux $\epsilon \approx 10^{-8} \text{ W/kg}$, yields a lifetime of $\approx 1 \text{ yr}$ which is indeed the scale separating a high frequency “ocean weather” (with $\beta > 1$) from a low frequency “macro-ocean weather” with $\beta < 1$ [*Lovejoy and Schertzer, 2012c*], see fig. 3.

This picture allows us to understand the weather/macroweather transition since it validates the use of the stochastic turbulence based Fractionally Integrated Flux model (FIF, i.e. cascades [*Schertzer and Lovejoy, 1987*]). The FIF model shows that whereas in the weather regime, fluctuations depend on interactions in both space and in time, at lower frequencies they become “quenched” so that only the temporal interactions are important and τ_w marks a “dimensional transition” [*Lovejoy and Schertzer, 2010*]. Physically, at scales $\Delta t < \tau_w$ the statistics depend on

structures with lifetimes Δt ; at scales $\Delta t > \tau_w$ they depend on the statistics of many planetary sized structures. In addition, the basic FIF model predicts [Lovejoy and Schertzer, 2012c] macroweather exponents typically in the range $0.2 < \beta_{mw} < 0.6$ (i.e. $-0.4 < H_{mw} < -0.2$).

4. Real space fluctuations and analyses

In spite of the now burgeoning evidence that the atmosphere's natural variability is scaling over various ranges, the idea has not received the attention it deserves and at least over decadal, centennial and millennial scales, the natural variability is still largely identified with quasi-periodic behaviours which – when present – have the advantage of being predictable (for examples of periodicities ranging from multidecadal to millennial scales see [Delworth et al., 1993], [Schlesinger and Ramankutty, 1994], [Mann and Park, 1994], [Mann et al., 1995], [Bond et al., 1997], [Isono et al., 2009]). An additional reason for this focus on quasi-periodic behavior is that whereas spectra are ideal for understanding periodic processes, they are not optimal for scaling processes. For these, the corresponding real space analyses are more straightforward to interpret; this is particularly true when comparing spectra from different data types with different resolutions. In this section we show how this works.

In order to understand the qualitatively different behaviors in fig. 2, consider fluctuations ΔT . In a scaling regime, these will change with scale as $\Delta T = \phi \Delta t^H$ where H is the fluctuation exponent (also called the “nonconservation” exponent; it is denoted “ H ” in honor of Edwin Hurst but in general – unless the process is Gaussian – it is not the

same as the Hurst exponent). φ is a controlling dynamical variable (e.g. a turbulent flux) whose mean $\langle \varphi \rangle$ is independent of the lag Δt (i.e. independent of the time scale). The behavior of the mean fluctuation is thus $\langle \Delta T \rangle \approx \Delta t^H$ so that if $H > 0$, on average fluctuations tend to grow with scale whereas if $H < 0$, they tend to decrease.

Although it is traditional (and often sufficient) to define fluctuations by absolute differences $\Delta T(\Delta t) = |T(t + \Delta t) - T(t)|$, for our purposes this is *not* sufficient. Instead we should use the absolute difference of the mean between t and $t + \Delta t/2$ and between $t + \Delta t/2$ and $t + \Delta t$. Technically, the latter corresponds to defining fluctuations using Haar wavelets rather than “poor man’s” wavelets (differences). While the latter is adequate for fluctuations increasing with scale (i.e. $H > 0$), mean absolute differences cannot decrease and so when $H < 0$, they do not correctly estimate fluctuations. The Haar fluctuation (which is useful for $-1 < H < 1$) is particularly easy to understand since (with proper “calibration”) in regions where $H > 0$, it can be made very close to the difference fluctuation and in regions where $H < 0$, it can be made close to another simple to interpret “tendency fluctuation”. While other techniques such as Detrended Fluctuation Analysis [*Peng et al.*, 1994], [*Kantelhardt et al.*, 2002; *Monetti et al.*, 2003] perform just as well for determining exponents, they have the disadvantage that their fluctuations are not at all easy to interpret (they are the standard deviations of the residues of polynomial regressions on the running sum of the original series; see [*Lovejoy and Schertzer*, 2012b]).

Once estimated, the variation of the fluctuations with scale can be quantified by using their statistics; the q^{th} order structure function $S_q(\Delta t)$ is particularly convenient:

$$S_q(\Delta t) = \langle \Delta T(\Delta t)^q \rangle \quad (1)$$

where “ $\langle, \rangle$ ” indicates ensemble averaging. In a scaling regime, $S_q(\Delta t)$ is a power law; $S_q(\Delta t) \approx \Delta t^{\xi(q)}$, where the exponent $\xi(q) = qH - K(q)$ where $K(1) = 0$. Since Gaussian processes have $K(q) = 0$, $K(q)$ characterizes the strong non Gaussian variability; the “intermittency”. In the macroweather regime $K(2)$ is typically small ($\approx 0.01-0.03$), so that the RMS variation $S_2(\Delta t)^{1/2}$ (denoted simply $S(\Delta t)$ below) has the exponent $\xi(2)/2 \approx \xi(1) = H$. In the climate regime this intermittency correction is a bit larger [Schmitt *et al.*, 1995] but the error in using this approximation (≈ 0.06) will be neglected. Note that since the spectrum is a second order statistic, we have the useful relationship $\beta = 1 + \xi(2) = 1 + 2H - K(2)$. When $K(2)$ is small, $\beta \approx 1 + 2H$ so that as mentioned earlier, $H > 0$, $H < 0$ corresponds to $\beta > 1$, $\beta < 1$ respectively.

When $S(\Delta t)$ is estimated for various in situ, reanalysis, multiproxy and paleo temperatures, one obtains fig. 3. The key points to note are a) the three qualitatively different regimes corresponding to the spectra in fig. 1: weather, (low frequency) macroweather and climate with $S(\Delta t)$ respectively increasing, decreasing and increasing again with scale ($H_w > 0$, $H_{mw} < 0$, $H_c > 0$). The transitions are at $\tau_w \approx 5 - 10$ days and $\tau_c \approx 30 - 100$ yrs: note that in the recent period τ_c is a bit lower (≈ 10 yrs) - see the 1880-2008 surface series - than in earlier periods (see the multiproxies with $\tau_c \approx 30 - 100$ yrs), this is because over the last century, anthropogenic forcing has

become dominant for scales greater than about 10 yrs, see [Lovejoy, 2013]. We can also glimpse a fourth low frequency “macro climate” regime for scales larger than $\tau_{mc} \approx 100$ *kyrs*, but this is outside our scope. b) the difference between the local and global fluctuations. c) the “glacial/interglacial window” corresponding to overall ± 3 to ± 5 *K* variations (i.e. $S(\Delta t) \approx 6, 10$ *K*) over scales with half periods of 30 – 50 *kyrs*; the curve must pass through the window in order to explain the glacial/interglacial transitions. Starting at $\tau_c \approx 10 - 30$ *yrs*, one can plausibly extrapolate the global surface and 20CR 700 *mb* $S(\Delta t)$ ’s using $H = 0.4$ ($\beta \approx 1.8$), all the way to the interglacial window (with nearly an identical S as in [Lovejoy and Schertzer, 1986]). Similarly, the local temperatures and multiproxies also seem to follow the same exponent with slightly different τ_c ’s and seem to extrapolate respectively a little above and below the window.

These statistics may seem arcane but their physical interpretation is pretty straightforward. In the weather regime, larger and larger fluctuations “live” for longer and longer times: the “eddy turnover time”. At any given time scale, the fluctuations are dominated by structures with corresponding spatial scales and this relationship holds up to structures of planetary scales with lifetimes ≈ 10 days. For periods longer than this, the statistics are dominated by averages of many planetary scale structures, and these fluctuations tend to cancel out: for example large temperature increases are typically followed (and partially cancelled) by corresponding decreases. The consequence is that in this macroweather regime, the average fluctuations diminish as the time scale increases. At some point – at around 30 years depending on geographic location and time (and as little as 10 years in the

recent period when anthropogenic variability is important), these weaker and weaker fluctuations - whose origin is in weather dynamics - become dominated by increasingly strong lower frequency processes. These not only include changing external solar, volcanic orbital or anthropogenic “forcings” – but quite likely also new and increasingly strong slow (internal) climate processes - or by a combination of the two: forcings with feedbacks. A relevant example of a slow dynamical process that is not currently fully incorporated into GCM’s is land-ice dynamics. The overall effect is that in the resulting climate regime, fluctuations tend to grow again with scale in an “unstable” manner, very similar to the way they grow in the weather regime.

5. Implications for climate modelling, prediction

Numerical weather models and reanalyses are qualitatively in good agreement with the weather / macroweather picture described above, although there are still some quantitative discrepancies in the values of the exponents, possibly due the hydrostatic approximation and other numerical issues ([*Stolle et al.*, 2009], [*Lovejoy and Schertzer*, 2011]). However, climate models (GCM’s) are essentially weather models with various additional couplings (with ocean, carbon cycle, land-use, sea ice and other models). It is therefore not surprising that control runs (i.e. with no “climate forcings”) generate macroweather (with $\beta_{mw} \approx 0.2$, $H_{mw} \approx -0.4$), and this apparently out to the extreme low frequency limit of the models (see the review and analyses in [*Lovejoy et al.*, 2012] as well as [*Blender et al.*, 2006], [*Rybski et al.*, 2008]).

Avoiding anthropogenic effects by considering the pre-1900 epoch, for GCM climate models, the key question is whether solar, volcanic, orbital or other climate forcings are sufficient to arrest the $H < 0$ decline in macroweather fluctuations and to create an $H > 0$ regime with sufficiently strong centennial, millennial variability to account for the background variability out to glacial- interglacial scales. Analysis of several last millennium simulations has found that at the moment, their low frequency variabilities are too weak [Lovejoy *et al.*, 2012].

To understand this weak variability, one can examine the scale dependence of fluctuations in the radiative forcings (ΔR_F) of several solar and volcanic reconstructions; they are generally scaling with $\Delta R_F \approx A \Delta t^H$ [Lovejoy and Schertzer, 2012a]. If $H_R \approx H_T \approx 0.4$, then scale *independent* amplification / feedback mechanisms would suffice. However it was often found that $H_R \approx -0.3$ implying that the forcings become weaker with scale - even though the response grows with scale. This suggests the need to introduce new slow mechanisms of internal variability. Such mechanisms must have broad spectra; this suggests that their dynamics involve nonlinearly interacting spatial degrees of freedom such as the land-ice dynamics mentioned above.

Whatever the ultimate source of the growing fluctuations in the $H > 0$ climate regime, a careful and complete characterization of the scaling in space as well as in time (including possible space-time anisotropies) allows for new stochastic methods for predicting the climate. The idea is to exploit the particularly low variability of the averages at scale τ_c . Since $\tau_c \approx 30$ yrs - i.e. the conventional but ad hoc “climate normal” period - this not only justifies the normal but allows averages of relevant

variables over it to define “climate states” and the changes at scales $\Delta t > \tau_c$ to define climate change (again, in the recent period, this defines the scale at which anthropogenic variability starts to dominate natural variability). Even without resolving the question of the dominant climate forcing and slow internal feedbacks, one could use the statistical properties of the climate states - the system’s “memory” implicit in the long range statistical correlations – combined with the growing data on past climate states in order to make stochastic climate forecasts.

Another attractive application of this scaling picture is that by quantifying the natural variability as a function of space-time scales, it opens up the possibility of convincingly distinguish natural and anthropogenic variability. This is possible because the stochastic scaling framework allows one to statistically test specific hypotheses about the probability that the atmosphere would naturally behave in the way that is observed, i.e. to formulate rigorous statistical tests of any trends or events against the null hypothesis. Only if the probabilities are low enough should the hypothesis that the observed changes are natural in origin be rejected. For example, using CO₂ as a surrogate for all anthropogenic effects and taking into both long range statistical dependencies and extreme “fat tailed” probabilities, [Lovejoy, 2013] found that the probability of the global warming since 1880 being due to natural variability can be rejected with more than 99% confidence. The scaling fluctuation approach thus allows quantitative (and hence convincing) answers to questions such as: how can the earth have prolonged periods of cooling in the midst of anthropogenic warming; or was this winter’s record mild temperature really evidence for anthropogenic influence? Finally, the systematic comparison of model

and natural variability in the preindustrial era is the best way to fully address the issue of “model uncertainty”, to assess the extent by which the models are missing important slow processes.

6. Conclusions

Contrary to [Bryson, 1997], we have argued that the climate is not accurately viewed as the statistics of fundamentally fast weather dynamics that are constrained by quasi fixed boundary conditions. The empirically substantiated picture is rather one of “unstable”, “wandering”, high frequency weather processes (i.e. $H > 0$) tending - at scales beyond 10 days or so and primarily due to the quenching of spatial degrees of freedom (intermediate frequency, low variability) to macroweather processes. These appear to be stable because positive and negative fluctuations tend to cancel out ($H < 0$).

True climate processes are “weather-like” ($H > 0$) and only emerge from macroweather at even lower frequencies, due to new slow internal climate processes coupled with external forcings (including in the recent period, anthropogenic forcings). These processes presumably include various nonlinear couplings with the fields that Bryson considered to be no more than “boundary conditions” so that “rather than ‘boundaries’ these become interactive mediums” [Pielke, 1998]. The theoretical/mathematical picture underpinning GCM based prediction beyond ≈ 30 years should thus be re-examined.

Yet, whatever the cause, it is an empirical fact that the emergent synergy of new processes yields fluctuations that on average again grow with scale in at least a roughly scaling manner and become dominant typically on time scales of 30 -100 years (somewhat less in the recent period) up to ≈ 100 *kyrs*.

Looked at another way, if the climate really *was* what you expected, then – since one usually expects averages - predicting the climate would be the relatively simple matter of determining the fixed climate normal. On the contrary, we have argued that from the stochastic point of view - and notwithstanding the vastly different time scales - that predicting natural climate change is very much like predicting the weather. This is because the climate at any time or place is the consequence of climate changes that are (qualitatively and quantitatively) unexpected in very much the same way that the weather is unexpected.

At a subjective level, from experience over the years, we all grow to expect certain stable patterns of macroweather (complicated by seasonal, and now by anthropogenic effects, but nevertheless recognizable from year to year) so that in daily discourse we may say “macroweather is what you expect, the weather is what you get”. However over generational scales - periods of 30 years - the macroweather we are accustomed to evolves due to climate change. Speaking to our children and grandchildren, the appropriate dictum would therefore be “macroweather is what you expect, the climate is what you get”.

6. Acknowledgements:

We would like to thank Roger Pielke, Gavin Schmidt, and Adrian Tuck for useful comments.

7. References

- AchutaRao, K., and K. R. Sperber (2006), ENSO simulation in coupled ocean-atmosphere models: are the current models better? , *Climate Dynamics*, 27, 1–15 doi: 10.1007/s00382-006-0119-7.
- Ashkenazy, Y., D. Baker, H. Gildor, and S. Havlin (2003), Nonlinearity and multifractality of climate change in the past 420,000 years, *Geophys. Res. Lett.*, 30, 2146 doi: 10.1029/2003GL018099.
- Blender, R., K. Fraedrich, and B. Hunt (2006), Millennial climate variability: GCM-simulation and Greenland ice cores, *Geophys. Res. Lett.*, , 33, , L04710 doi: doi:10.1029/2005GL024919.
- Bond, G., W. Showers, M. Cheseby, R. Lotti, P. Almasi, P. deMenocal, P. Priori, H. Cullen, I. Hajdes, and G. Bonani (1997), A pervasive millennial-scale climate cycle in the North Atlantic: The Holocene and late glacial record, *Science* 278, 1257-1266.
- Bryson, R. A. (1997), The Paradigm of Climatology: An Essay, *Bull. Amer. Meteor. Soc.* , 78, 450-456.
- Charney, J. G. (1971), Geostrophic Turbulence, *J. Atmos. Sci.*, 28, 1087.
- Committee on Radiative Forcing Effects on Climate, N. R. C. (2005), *Radiative Forcing of Climate Change: Expanding the Concept and Addressing Uncertainties*, 224 pp., National Acad. press.
- Delworth, T., S. Manabe, and R. J. Stoufer (1993), Interdecadal variations of the thermocline circulation in a coupled ocean-atmosphere model, *J. of Climate*, 6, 1993-2011.
- Ditlevsen, P. D., H. Svensmark, and S. Johson (1996), Contrasting atmospheric and climate dynamics of the last-glacial and Holocene periods, *Nature*, 379, 810-812.
- Eichner, J. F., E. Koscielny-Bunde, A. Bunde, S. Havlin, and H.-J. Schellnhuber (2003), Power-law persistence and trends in the atmosphere: A detailed study of long temperature records, *Phys. Rev. E*, 68, 046133-046131-046135 doi: 10.1103/PhysRevE.68.046133.
- Fraedrich, K., and K. Blender (2003), Scaling of Atmosphere and Ocean Temperature Correlations in Observations and Climate Models, *Phys. Rev. Lett.*, 90, 108501-108504
- Fraedrich, K., R. Blender, and X. Zhu (2009), Continuum Climate Variability: Long-Term Memory, Scaling, and 1/f-Noise, , *International Journal of Modern Physics B*, 23, 5403-5416.
- Heinlein, R. A. (1973), *Time Enough for Love*, 605 pp., G. P. Putnam's Sons, New York.
- Huang, S. (2004), Merging Information from Different Resources for New Insights into Climate Change in the Past and Future, *Geophys. Res. Lett.* , 31, L13205 doi: doi : 10.1029/2004 GL019781.
- Huschke, R. E. (Ed.) (1959), *Glossary of Meteorology*, 638 pp. pp.
- Huybers, P., and W. Curry (2006), Links between annual, Milankovitch and continuum temperature variability, *Nature*, 441, 329-332 doi: 10.1038/nature04745.

- 545 Isono, D., M. Yamamoto, T. Irino, T. Oba, M. Murayama, T. Nakamura, and H.
 546 Kawahata (2009), The 1500-year climate oscillation in the midlatitude North
 547 Pacific during the Holocene, *Geology* 37(591-594).
- 548 Kantelhardt, J. W., S. A. Zschechner, K. Koscielny-Bunde, S. Havlin, A. Bunde, and H.
 549 E. Stanley (2002), Multifractal detrended fluctuation analysis of nonstationary time
 550 series, *Physica A*, 316, 87-114.
- 551 Kolesnikov, V. N., and A. S. Monin (1965), Spectra of meteorological field fluctuations,
 552 *Izvestiya, Atmospheric and Oceanic Physics*, 1, 653-669.
- 553 Koscielny-Bunde, E., A. Bunde, S. Havlin, H. E. Roman, Y. Goldreich, and H. J.
 554 Schellnhuber (1998), Indication of a universal persistence law governing
 555 atmospheric variability, *Phys. Rev. Lett.*, 81, 729.
- 556 Kraichnan, R. H. (1967), Inertial ranges in two-dimensional turbulence, *Physics of Fluids*,
 557 10, 1417-1423.
- 558 Lamb, H. H. (1972), *Climate: Past, Present, and Future. Vol. 1, Fundamentals and*
 559 *Climate Now*, Methuen and Co.
- 560 Lanfredi, M., T. Simoniello, V. Cuomo, and M. Macchiato (2009), Discriminating low
 561 frequency components from long range persistent fluctuations in daily atmospheric
 562 temperature variability, *Atmos. Chem. Phys.*, 9, 4537-4544.
- 563 Lennartz, S., and A. Bunde (2009), Trend evaluation in records with long term memory:
 564 Application to global warming, *Geophys. Res. Lett.*, 36, L16706 doi:
 565 doi:10.1029/2009GL039516.
- 566 Lindborg, E., K. K. Tung, G. D. Nastrom, J. Y. N. Cho, and K. S. Gage (2010a),
 567 Comment on "Reinterpreting aircraft measurement in anisotropic scaling turbulence"
 568 by Lovejoy et al., *Atmos. Chem. Phys.*, 10, 1401-1402.
- 569 Lindborg, E., K. K. Tung, G. D. Nastrom, J. Y. N. Cho, and K. S. Gage (2010b),
 570 Interactive comment on "Comment on "Reinterpreting aircraft measurements in
 571 anisotropic scaling turbulence" by Lovejoy et al. (2009)", *Atmos. Chem. Phys.*
 572 *Discuss.*, 9 C9797-C9798.
- 573 Ljungqvist, F. C. (2010), A new reconstruction of temperature variability in the extra -
 574 tropical Northern Hemisphere during the last two millennia, *Geografiska Annaler:*
 575 *Physical Geography*, 92 A(3), 339 - 351 doi: DOI : 10.1111/j .1468 -
 576 0459.2010 .00399.x.
- 577 Lorenz, E. N. (1995), *Climate is what you expect*, edited, p. 55pp,
 578 aps4.mit.edu/research/Lorenz/publications.htm, (available, 16 May, 2012).
- 579 Lovejoy, S. (2013), Anthropogenic warming quantified by scaling fluctuation analysis
 580 *Nature Clim Change*, (submitted).
- 581 Lovejoy, S., and D. Schertzer (1984), 40 000 years of scaling in climatological
 582 temperatures, *Meteor. Sci. and Tech.*, 1, 51-54.
- 583 Lovejoy, S., and D. Schertzer (1986), Scale invariance in climatological temperatures and
 584 the spectral plateau, *Annales Geophysicae*, 4B, 401-410.
- 585 Lovejoy, S., and D. Schertzer (2010), Towards a new synthesis for atmospheric
 586 dynamics: space-time cascades, *Atmos. Res.*, 96, pp. 1-52 doi: doi:
 587 10.1016/j.atmosres.2010.01.004.
- 588 Lovejoy, S., and D. Schertzer (2011), Space-time cascades and the scaling of ECMWF
 589 reanalyses: fluxes and fields, *J. Geophys. Res.*, 116 doi: 10.1029/2011JD015654.

- 590 Lovejoy, S., and D. Schertzer (2012a), Stochastic and scaling climate sensitivities: solar,
 591 volcanic and orbital forcings, *Geophys. Res. Lett.* , 39, L11702 doi:
 592 doi:10.1029/2012GL051871.
- 593 Lovejoy, S., and D. Schertzer (2012b), Haar wavelets, fluctuations and structure
 594 functions: convenient choices for geophysics, *Nonlinear Proc. Geophys.* , 19, 1-14
 595 doi: 10.5194/npg-19-1-2012.
- 596 Lovejoy, S., and D. Schertzer (2012c), Low frequency weather and the emergence of the
 597 Climate, in *Extreme Events and Natural Hazards: The Complexity Perspective*,
 598 edited by A. S. Sharma, A. Bunde, D. Baker and V. P. Dimri, pp. 231-254, AGU
 599 monographs.
- 600 Lovejoy, S., and D. Schertzer (2013), *The Weather and Climate: Emergent Laws and*
 601 *Multifractal Cascades*, 480 pp., Cambridge University Press, Cambridge.
- 602 Lovejoy, S., D. Schertzer, and A. F. Tuck (2010), Why anisotropic turbulence matters:
 603 another reply to E. Lindborg, *Atmos. Chem and Physics Disc.*, 10, C4689-C4697.
- 604 Lovejoy, S., D. Schertzer, and D. Varon (2012), Do GCM's predict the climate.... or
 605 macroweather?, *Earth Syst. Dynam. Discuss.*, 3, , 1259-1286 doi: 10.5194/esdd-3-
 606 1259-2012.
- 607 Lovejoy, S., A. F. Tuck, S. J. Hovde, and D. Schertzer (2007), Is isotropic turbulence
 608 relevant in the atmosphere?, *Geophys. Res. Lett.*, doi:10.1029/2007GL029359,
 609 L14802.
- 610 Lovejoy, S., A. F. Tuck, D. Schertzer, and S. J. Hovde (2009), Reinterpreting aircraft
 611 measurements in anisotropic scaling turbulence, *Atmos. Chem. and Phys.* , 9, 1-19.
- 612 Mann, M. E., and J. Park (1994), Global scale modes of surface temperature variability
 613 on interannual to century timescales, *J. Geophys. Resear.*, 99, 819-825.
- 614 Mann, M. E., J. Park, and R. S. Bradley (1995), Global interdecadal and century scale
 615 climate oscillations during the past five centuries, *Nature*, 378, 268-270.
- 616 Mitchell, J. M. (1976), An overview of climatic variability and its causal mechanisms,
 617 *Quaternary Res.*, 6, 481-493.
- 618 Moberg, A., D. M. Sonnechkin, K. Holmgren, and N. M. Datsenko (2005), Highly
 619 variable Northern Hemisphere temperatures reconstructed from low- and high -
 620 resolution proxy data, *Nature*, 433(7026), 613-617.
- 621 Monetti, R. A., S. Havlin, and A. Bunde (2003), Long-term persistence in the sea surface
 622 temperature fluctuations, *Physica A*, 320, 581-589.
- 623 Monin, A. S. (1972), *Weather forecasting as a problem in physics*, MIT press, Boston Ma.
- 624 Palmer, T. (2005), Global warming in a nonlinear climate – Can we be sure? ,
 625 *Europhysics news March/April 2005*, 42-46 doi: doi.org/10.1051/epn:2005202.
- 626 Palmer, T. N. (2012), Towards the probabilistic Earth-system simulator: a vision for the
 627 future of climate and weather prediction, *Q.J.R. Meteorol. Soc.*, in press.
- 628 Palmer, T. N., F. J. Doblas-Reyes, A. Weisheimer, and M. J. Rodwell (2008), Toward
 629 Seamless Prediction: Calibration of Climate Change Projections Using Seasonal
 630 Forecasts. , *Bull. Amer. Meteor. Soc.* , 89, 459–470 doi: doi.org/10.1175/BAMS-
 631 89-4-459.
- 632 Panofsky, H. A. (1969), The spectrum of temperature, *J. of Radio Science*, 4, 1101-1109.
- 633 Pelletier, J., D. (1998), The power spectral density of atmospheric temperature from
 634 scales of 10^{-2} to 10^6 yr , *EPSL*, 158, 157-164.

- Peng, C.-K., S. V. Buldyrev, S. Havlin, M. Simons, H. E. Stanley, and A. L. Goldberger (1994), Mosaic organisation of DNA nucleotides, *Phys. Rev. E*, **49**, 1685-1689.
- Pielke, R. (1998), Climate prediction as an initial value problem, *Bull. of the Amer. Meteor. Soc.*, **79**, 2743-2746.
- Pielke, R. A. S., R. Wilby, D. Niyogi, F. Hossain, K. Dairuku, J. Adegoke, G. Kallos, T. Seastedt, and K. Suding (2012), Dealing with Complexity and Extreme Events Using a Bottom-up, Resource-based Vulnerability Perspective, in *Complexity and Extreme Events in Geosciences*, edited by A. S. Sharma, A. Bunde, D. Baker and V. P. Dimri, AGU.
- Pinel, J., S. Lovejoy, D. Schertzer, and A. F. Tuck (2012), Joint horizontal - vertical anisotropic scaling, isobaric and isoheight wind statistics from aircraft data, *Geophys. Res. Lett.*, **39**, L11803 doi: 10.1029/2012GL051698.
- Radkevitch, A., S. Lovejoy, K. B. Strawbridge, D. Schertzer, and M. Lilley (2008), Scaling turbulent atmospheric stratification, Part III: empirical study of Space-time stratification of passive scalars using lidar data, *Quart. J. Roy. Meteor. Soc.*, DOI: 10.1002/qj.1203.
- Rybski, D., A. Bunde, and H. von Storch (2008), Long-term memory in 1000- year simulated temperature records, *J. Geophys. Resear.*, **113**, D02106-02101, D02106-02109 doi: doi:10.1029/2007JD008568.
- Schertzer, D., and S. Lovejoy (1985), The dimension and intermittency of atmospheric dynamics, in *Turbulent Shear Flow 4*, edited by B. Launder, pp. 7-33, Springer-Verlag.
- Schertzer, D., and S. Lovejoy (1987), Physical modeling and Analysis of Rain and Clouds by Anisotropic Scaling of Multiplicative Processes, *Journal of Geophysical Research*, **92**, 9693-9714.
- Schertzer, D., I. Tchiguirinskaia, S. Lovejoy, and A. F. Tuck (2011), Quasi-geostrophic turbulence and generalized scale invariance, a theoretical reply to Lindborg, *Atmos. Chem. and Physics Discussion*, **11**, 3301-3320.
- Schertzer, D., I. Tchiguirinskaia, S. Lovejoy, and A. F. Tuck (2012), Quasi-geostrophic turbulence and generalized scale invariance, a theoretical reply, *Atmos. Chem. Phys.*, **12**, 327-336 doi: 10.5194/acp-12-327-2012.
- Schlesinger, M. E., and N. Ramankutty (1994), An Oscillation in the Global Climate System of Period 65-70 Years, *Nature*, **367**, 723-726.
- Schmitt, F., S. Lovejoy, and D. Schertzer (1995), Multifractal analysis of the Greenland Ice-core project climate data., *Geophys. Res. Lett.*, **22**, 1689-1692.
- Shackleton, N. J., and J. Imbrie (1990), The $\delta^{18}\text{O}$ spectrum of oceanic deep water over a five-decade band, *Climatic Change*, **16**, 217-230.
- Stolle, J., S. Lovejoy, and D. Schertzer (2009), The stochastic cascade structure of deterministic numerical models of the atmosphere, *Nonlin. Proc. in Geophys.*, **16**, 1-15.
- Talkner, P., and R. O. Weber (2000), Power spectrum and detrended fluctuation analysis: Application to daily temperatures, *Phys. Rev. E*, **62**, 150-160.
- Vallis, G. (2010), Mechanisms of climate variability from years to decades, in *Stochastic Physics and Climate Modelling*, edited by P. W. T. Palmer, pp. 1-34, Cambridge University Press, Cambridge.

- 680 Van der Hoven, I. (1957), Power spectrum of horizontal wind speed in the frequency
681 range from .0007 to 900 cycles per hour, *Journal of Meteorology*, *14*, 160-164.
682 Wunsch, C. (2003), The spectral energy description of climate change including the 100
683 ky energy, *Climate Dynamics*, *20*, 353-363.
684 Yano, J. (2009), Interactive comment on “Reinterpreting aircraft measurements in
685 anisotropic scaling turbulence” by S. Lovejoy et al., *Atmos. Chem. Phys. Discuss.*, ,
686 9, S162–S166 doi: <http://www.atmos-chem-phys-discuss.net/9/S162/2009/>.
687
688

689

Figures and Captions:

Fig. 1: A composite temperature spectrum. Left: the GRIP (Summit) ice core $\delta^{18}\text{O}$ (a temperature proxy, low resolution (55 cm in depth back 240 kyrs) along with the first 91 kyrs at high resolution. Right: the spectrum of the (mean) 75°N of the 20th Century Reanalysis (20CR) temperature spectrum, at 6 hour resolution, from 1871-2008, at 700 mb. The overlap (from 10 – 138 yr scales) is used for calibrating the former (moving them vertically on the log-log plot). All spectra are averaged over logarithmically spaced bins, ten per order of magnitude in frequency. Three regimes are shown corresponding to the weather regime with $\beta_w = 2$ (the diurnal variation and subharmonic at 12 hours are visible at the extreme right). The central macroweather “plateau” is shown along with the theoretically predicted $\beta_{mw} = 0.2 - 0.4$ regime. Finally, at longer time scales (the left), a

703 new scaling climate regime with exponent $\beta_c \approx 1.4$ continues to about 100 *kyrs*.

704 Reproduced from [Lovejoy and Schertzer, 2012c].

705

706

707 **Fig. 2: Dynamics and types of scaling variability:** A visual comparison displaying
 708 representative temperature series from weather, macroweather and climate ($H \approx$
 709 0.4, -0.4, 0.4, bottom to top respectively). To make the comparison as fair as possible,
 710 in each case, the sample is 720 points long and each series has its mean removed
 711 and is normalized by its standard deviation (4.49 K, 2.59 K, 1.39 K, respectively), the
 712 two upper series have been displaced in the vertical by four units for clarity. The
 713 resolutions are 1 hour, 20 days and 100 *yrs*, the data are from a weather station in
 714 Lander Wyoming (detrended daily, annually), the 20th Century reanalysis and the

715 Vostok Antarctic station. Note the similarity between the type of variability in the
716 weather and climate regimes (reflected in their scaling exponents although the H
717 exponent is only a partial characterization).

718

719

720 **Fig. 3: Empirical RMS temperature fluctuations ($S(\Delta t)$), local scale analyses:** On
 721 the left top we show grid point scale ($2^\circ \times 2^\circ$) daily scale fluctuations globally
 722 averaged along with reference slope $\xi(2)/2 = -0.4 \approx H$ (20CR, 700 mb). For
 723 comparison, the results for 50 simulations of Ornstein-Uhlenbeck (OU) processes are
 724 also given using simulations with a characteristic time of 3 days. The theoretical
 725 asymptotic slopes (0.5, -0.5) are added to show their convergence to theory. Just below
 726 this, we show the monthly NOAA CDC Sea Surface Temperature curve (5° resolution,
 727 from 1900-2000); the transition from $\xi(2)/2 \approx 0.4$ to ≈ -0.2 occurs at $\tau_{ow} \approx 1 \text{ yr}$. On the
 728 lower left, we see at daily resolution, the corresponding globally averaged structure
 729 function.

730 **Globally averaged series:** The same 20CR data but globally averaged (brown), The

average of the three in situ surface series (NOAA NCDC, NASA GISS, CRU, red) as well as the average of three post 2003 multiproxy structure functions; [*Huang, 2004*], [*Ljundqvist, 2010; Moberg et al., 2005*], (see [*Lovejoy and Schertzer, 2012c*]).

Paleotemperatures: At the right we show analysis of the EPICA Antarctic series interpolated to 50 year resolution over ≈ 800 *kyrs*. Also shown is the interglacial “window”.

The reference slopes are $\xi(2)/2 = -0.4, -0.2$ or $+0.4$; these correspond to spectral exponents $\beta = 1 + \xi(2) = 0.2, 0.6, 1.8$ respectively; a flat curve in the above corresponds to $\beta = 1$.

Figure Captions:

Fig. 1: **A composite temperature spectrum.** Left: the GRIP (Summit) ice core $\delta^{18}\text{O}$ (a temperature proxy, low resolution (55 cm in depth back 240 kyrs) along with the first 91 kyrs at high resolution. Right: the spectrum of the (mean) 75°N of the 20th Century Reanalysis (20CR) temperature spectrum, at 6 hour resolution, from 1871-2008, at 700 mb. The overlap (from 10 – 138 yr scales) is used for calibrating the former (moving them vertically on the log-log plot). All spectra are averaged over logarithmically spaced bins, ten per order of magnitude in frequency. Three regimes are shown corresponding to the weather regime with $\beta_w = 2$ (the diurnal variation and subharmonic at 12 hours are visible at the extreme right). The central macroweather “plateau” is shown along with the theoretically predicted $\beta_{mw} = 0.2 - 0.4$ regime. Finally, at longer time scales (the left), a new scaling climate regime with exponent $\beta_c \approx 1.4$ continues to about 100 kyrs. Reproduced from [Lovejoy and Schertzer, 2012c].

Fig. 2: **Dynamics and types of scaling variability:** A visual comparison displaying representative temperature series from weather, macroweather and climate ($H \approx 0.4, -0.4, 0.4$, bottom to top respectively). To make the comparison as fair as possible, in each case, the sample is 720 points long and each series has its mean removed and is normalized by its standard deviation (4.49 K, 2.59 K, 1.39 K, respectively), the two upper series have been displaced in the vertical by four units for clarity. The resolutions are 1 hour, 20 days and 100 yrs, the data are from a weather station in Lander Wyoming (detrended daily, annually), the 20th Century reanalysis and the Vostok Antarctic station. Note the similarity between the type of variability in the

weather and climate regimes (reflected in their scaling exponents although the H exponent is only a partial characterization).

Fig. 3: Empirical RMS temperature fluctuations ($S(\Delta t)$), local scale analyses: On the left top we show grid point scale ($2^\circ \times 2^\circ$) daily scale fluctuations globally averaged along with reference slope $\xi(2)/2 = -0.4 \approx H$ (20CR, 700 mb). For comparison, the results for 50 simulations of Ornstein-Uhlenbeck (OU) processes are also given using simulations with a characteristic time of 3 days. The theoretical asymptotic slopes (0.5, -0.5) are added to show their convergence to theory. Just below this, we show the monthly NOAA CDC Sea Surface Temperature curve (5° resolution, from 1900-2000); the transition from $\xi(2)/2 \approx 0.4$ to ≈ -0.2 occurs at $\tau_{ow} \approx 1$ yr. On the lower left, we see at daily resolution, the corresponding globally averaged structure function.

Globally averaged series: The same 20CR data but globally averaged (brown), The average of the three in situ surface series (NOAA NCDC, NASA GISS, CRU, red) as well as the average of three post 2003 multiproxy structure functions; [Huang, 2004], [Ljungqvist, 2010; Moberg et al., 2005], (see [Lovejoy and Schertzer, 2012c]).

Paleotemperatures: At the right we show analysis of the EPICA Antarctic series interpolated to 50 year resolution over ≈ 800 kyrs. Also shown is the interglacial “window”.

The reference slopes are $\xi(2)/2 = -0.4, -0.2$ or $+0.4$; these correspond to spectral exponents $\beta = 1 + \xi(2) = 0.2, 0.6, 1.8$ respectively; a flat curve in the above corresponds to $\beta = 1$.

786